

Lusternik-Schnirelmann category of non-simply connected compact simple Lie groups

Norio Iwase^{a,1} Mamoru Mimura^b Tetsu Nishimoto^c

^a*Faculty of Mathematics, Kyushu University, Ropponmatsu Fukuoka 810-8560,
Japan*

^b*Department of Mathematics, Faculty of Science, Okayama University, 3-1
Tsushima-naka, Okayama 700-8530, Japan*

^c*Department of Welfare Business, Kinki Welfare University, Fukusaki-cho, Hyogo
679-2217, Japan*

Abstract

Let $F \hookrightarrow X \rightarrow B$ be a fibre bundle with structure group G , where B is $(d-1)$ -connected and of finite dimension, $d \geq 1$. We prove that the strong L-S category of X is less than or equal to $m + \frac{\dim B}{d}$, if F has a cone decomposition of length m under a compatibility condition with the action of G on F . This gives a consistent prospect to determine the L-S category of non-simply connected Lie groups. For example, we obtain $\text{cat}(PU(n)) \leq 3(n-1)$ for all $n \geq 1$, which might be best possible, since we have $\text{cat}(PU(p^r)) = 3(p^r-1)$ for any prime p and $r \geq 1$. Similarly, we obtain the L-S category of $SO(n)$ for $n \leq 9$ and $PO(8)$. We remark that all the above Lie groups satisfy the Ganea conjecture on L-S category.

Key words: Lusternik-Schnirelmann category; cone decomposition; Lie group; Ganea conjecture

1991 MSC: Primary 55M30, Secondary 22E20, 57N60

Email addresses: iwase@math.kyushu-u.ac.jp (Norio Iwase),
mimura@math.okayama-u.ac.jp (Mamoru Mimura), nishimoto@kinwu.ac.jp
(Tetsu Nishimoto).

¹ The first named author is supported by the Grant-in-Aid for Scientific Research #14654016 from Japan Society for the Promotion of Science.

1 Introduction

The Lusternik-Schnirelmann category $\text{cat}(X)$, L-S category for short, is the least integer m such that there is a covering of X by $(m+1)$ open subsets each of which is contractible in X .

Ganea [5] introduced a stronger notion of L-S category, $\text{Cat}(X)$, which is equal to the cone-length, that is, the least integer m such that there is a set of cofibre sequences $\{A_i \rightarrow X_{i-1} \hookrightarrow X_i\}_{1 \leq i \leq m}$ with $X_0 = \{*\}$ and X_m homotopy equivalent to X .

The weak L-S category $w\text{cat}(X)$ is the least integer m such that the reduced diagonal map $\bar{\Delta}^{m+1} : X \rightarrow \bigwedge^{m+1} X$ is trivial where $\bigwedge^{m+1} X$ is the smash product. The stabilised version of the invariant $w\text{cat}(X)$ is given as the least integer m such that the reduced diagonal map $\bar{\Delta}^{m+1} : X \rightarrow \bigwedge^{m+1} X$ is *stably* trivial. Let us denote it by $\text{cup}(X)$, the *cup-length* of X .

In 1971, Ganea [6] posed 15 problems on L-S category and its related topics: Computation of L-S category for various manifolds is given as the first problem and the second problem is known as the Ganea conjecture on L-S category. These problems especially the first two problems have attracted many authors such as James and Singhof [15], [28], [25], [26], [27], [16], Gómez-Larrañaga and González-Acuña [7], Montejano [18], Oprea and Rudyak [20], [21], [19] and the authors [10], [11], [12], [13], [14]. In [11,12], the first author gave a counter example as a manifold to the Ganea conjecture on L-S category.

Especially for L-S category of compact connected simple Lie groups, the followings have already been known:

$$\begin{aligned} \text{cat}(\text{Sp}(1)) &= \text{cat}(\text{SU}(2)) = \text{cat}(\text{Spin}(3)) = 1, \\ \text{cat}(\text{SU}(3)) &= 2, \quad \text{cat}(\text{SO}(3)) = 3, \end{aligned}$$

since $\text{Sp}(1) = \text{SU}(2) = \text{Spin}(3) = S^3$, $\text{SU}(3) = \Sigma \mathbb{C}P^2 \cup e^8$ and $\text{SO}(3) = \mathbb{R}P^3$. Schweitzer [24] showed

$$\text{cat}(\text{Sp}(2)) = 3$$

using functional cohomology operations. Singhof [25,27] showed

$$\begin{aligned} \text{cat}(\text{SU}(n)) &= n-1, \\ \text{cat}(\text{Sp}(n)) &\geq n+1, \quad \text{if } n \geq 2. \end{aligned}$$

Also we know

$$\text{cat}(\text{G}_2) = 4$$

by [15] (see [13]). James and Singhof [16] showed

$$\text{cat}(\text{SO}(5)) = 8.$$

The first and second authors [13] and Fernández-Suárez, Gómez-Tato, Strom and Tanré [4] proved

$$\begin{aligned}\text{cat}(\text{Sp}(3)) &= 5, \\ \text{cat}(\text{Sp}(n)) &\geq n + 2 \quad \text{if } n \geq 3,\end{aligned}$$

by showing the reduced diagonal $\bar{\Delta}^5$ is given by the Toda bracket $\{\eta, \nu, \eta\} = \nu^2$. The authors [14] showed

$$\text{cat}(\text{Spin}(7)) = 5, \quad \text{cat}(\text{Spin}(8)) = 6$$

using explicit cone decompositions of $\text{Spin}(7)$ and $\text{SU}(4)$. Then the Ganea conjecture on L-S category holds for all these Lie groups, since the L-S and the strong L-S categories are equal to the cup-length:

Fact 1.1 *If $\text{cat}(X) = \text{cup } X$, then the Ganea conjecture on L-S category holds for X , i.e., $\text{cat}(X \times S^n) = \text{cat}(X) + 1$ for all $n \geq 1$.*

In fact, we have $\text{cup}(X \times S^n) = \text{cup}(X) + 1$ in general.

For any multiplicative cohomology theory h , we define $\text{cup}(X; h)$, the *cup-length* with respect to h , by the least integer m such that $u_0 \cdots u_m = 0$ for any $m+1$ elements $u_i \in \tilde{h}^*(X)$. When h is the ordinary cohomology theory with coefficient ring R , $\text{cup}(X; h)$ is often denoted as $\text{cup}(X; R)$.

Theorem 1.2 *For any CW-complex X we have*

$$\text{cup}(X) = \max\{\text{cup}(X; h) \mid h \text{ is any multiplicative cohomology theory}\}.$$

Proof. It is easy to see that $\text{cup}(X) \geq \text{cup}(X; h)$, and hence we have $\text{cup}(X) \geq \max\{\text{cup}(X; h) \mid h \text{ is any multiplicative cohomology theory}\}$. Thus we must show

$$\text{cup}(X) \leq \max\{\text{cup}(X; h) \mid h \text{ is any multiplicative cohomology theory}\}.$$

Let $m = \max\{\text{cup}(X; h) \mid h \text{ is any multiplicative cohomology theory}\}$ and h_X be the multiplicative cohomology theory represented by the following wedge sum of iterated smash products of suspension spectrum $\Sigma^\infty X$:

$$S^0 \vee \Sigma^\infty X \vee \Sigma^\infty \wedge^2 X \vee \cdots \vee \Sigma^\infty \wedge^i X \vee \cdots .$$

Let $\iota \in \tilde{h}_X^*(X)$ be the element which is represented by the inclusion map into the second factor $\Sigma^\infty X$ of the above wedge sum. Then by the definition of the cup-length, we have $\iota^{m+1} = 0$ which is represented by the reduced diagonal map $\bar{\Delta}^{m+1} : X \rightarrow \wedge^{m+1} X$ in the $(m+2)$ -nd factor $\Sigma^\infty \wedge^{m+1} X$ of the above wedge sum. Hence we have $\text{cup}(X) \leq m$ the desired inequality. Thus we obtain the result. $\square$

Let $P^m(\Omega X)$ be the m -th projective space, in the sense of Stasheff [29], such that there is a homotopy equivalence $P^\infty(\Omega X) \simeq X$. The following theorem is obtained by Ganea (see also [10] and Sakai [23]).

Theorem 1.3 (Ganea [5]) $\text{cat}(X) \leq m$ if and only if there is a map $\sigma : X \rightarrow P^m(\Omega X)$ such that $e_m^X \circ \sigma \sim 1_X$, where $e_m^X : P^m(\Omega X) \hookrightarrow P^\infty(\Omega X) \simeq X$.

Using this, Rudyak [21,22] introduced a stable L-S category, $\text{rcat}(X)$, which is the least integer m such that there is a stable map $\sigma : X \rightarrow P^m(\Omega X)$ satisfying $e_m^X \circ \sigma \sim 1_X$, another stabilised version of L-S category.

Rudyak [20] [21] and Strom [30] introduced the following invariant to calculate $\text{rcat}(X)$: Let $\text{wgt}(X; h)$ be the least integer m such that the homomorphism $(e_m^X)^* : \tilde{h}^*(X) \rightarrow \tilde{h}^*(P^m(\Omega X))$ is injective for any cohomology theory h . When h is the ordinary cohomology theory with coefficient ring R , $\text{wgt}(X; h)$ is often denoted as $\text{wgt}(X; R)$.

Since a product of any $m+1$ elements of $\tilde{h}^*(P^m(\Omega X))$ is trivial, we have $\text{cup}(X; h) \leq \text{wgt}(X; h)$ for any multiplicative cohomology theory h . Hence we have $\text{cup}(X) \leq \text{wgt}(X)$, where we denote $\text{wgt}(X) = \max\{\text{wgt}(X; h) \mid h \text{ is any cohomology theory}\}$.

Remark 1.4 For any ring R , we know $\text{cup}(\text{Sp}(2); R) = \text{wgt}(\text{Sp}(2); R) = 2 < 3 = \text{cat}(\text{Sp}(2))$. But an easy calculation of algebra structure of $KO^*(\text{Sp}(2))$ yields $\text{cup}(\text{Sp}(2); KO) = \text{wgt}(\text{Sp}(2); KO) = 3 = \text{cat}(\text{Sp}(2))$.

The following theorem is due to Rudyak [21,22], although we do not know the precise relation between $\text{wcat}(X)$ and $\text{rcat}(X)$.

Theorem 1.5 For any CW complex X , we have

$$\text{rcat}(X) = \text{wgt } X$$

and hence we have the following relations among categories:

$$\text{cup}(X) \leq \text{wcat}(X), \text{rcat}(X) \leq \text{cat}(X) \leq \text{Cat}(X).$$

Using this stabilised version of L-S category, we have the following theorem.

Theorem 1.6 (Rudyak [21,22]) If $\text{cat}(X) = \text{rcat}(X)$, then the Ganea conjecture on L-S category holds for X .

In fact, we have $\text{rcat}(X \times S^n) = \text{rcat}(X) + 1$ in general ([21,22]).

2 Main results

From now on, we work in the category of connected CW-complexes and continuous maps. We denote by $Z^{(k)}$ the k -skeleton of a CW complex Z .

Theorem 2.1 (James [15], Ganea [5]) *Let X be a $(d-1)$ -connected space of finite dimension. Then $\text{cat}(X) \leq \text{Cat}(X) \leq \lceil \frac{\dim(X)}{d} \rceil$, where $[a]$ denotes the biggest integer $\leq a$.*

In this paper, we extend this for a total space of a fibre bundle, to determine L-S categories of $\text{SO}(n)$ for $n \leq 9$, $\text{PO}(8)$ and $\text{PU}(p^r)$ (and the other quotient groups of $\text{SU}(p^r)$), which also gives an alternative proof of a result due to James and Singhof [16] on $\text{SO}(5)$.

We assume that B is a $(d-1)$ -connected finite dimensional CW complex ($d \geq 1$), whose cells are concentrated in dimensions $0, 1, \dots, s \pmod d$ for some s , ($0 \leq s \leq d-1$). Let $F \hookrightarrow X \rightarrow B$ be a fibre bundle with structure group G , a compact Lie group. Then we have the associated principal bundle $G \hookrightarrow E \xrightarrow{\pi} B$ with G -action $\psi : G \times F \rightarrow F$ on F and hence $X = E \times_G F$.

Let $K_i \xrightarrow{\rho_i} F_{i-1} \hookrightarrow F_i$, ($1 \leq i \leq m$) be m cofibre sequences with $F_0 = \{*\}$ and F_m homotopy equivalent to F . We consider the following compatibility condition of the above cone decomposition of F and the action of G on F .

Assumption 1 $\psi|_{G^{(d \cdot (i+1) + s - 1)} \times F_j} : G^{(d \cdot (i+1) + s - 1)} \times F_j \rightarrow F$ is compressible into F_{i+j} , $0 \leq i, j \leq i+j \leq m$.

Remark 2.2 (1) *Let $F = G$ and $X = E$ be the total space of a principal bundle over a path-connected space B and $d = 1$. Then any cone decomposition of F such that $F_i = F^{(n_i)}$ with $0 < n_1 < n_2 < \dots < n_m = \dim(F)$ satisfies Assumption 1 with $s = 0$.*

(2) *Let $F \hookrightarrow X \rightarrow B$ be a trivial bundle. Then any cone decomposition of F satisfies the compatibility Assumption 1 with $s = d-1$.*

Our main result is stated as follows:

Theorem 2.3 *Let B be a $(d-1)$ -connected finite dimensional CW complex ($d \geq 1$), whose cells are concentrated in dimensions $0, 1, \dots, s \pmod d$ for some s , $0 \leq s \leq d-1$. Let $F \hookrightarrow X \rightarrow B$ be a fibre bundle with fibre F whose structure group is a compact Lie group G . If F has a cone decomposition with the compatibility Assumption 1 for d , then $\text{Cat}(X) \leq m + \lceil \frac{\dim B}{d} \rceil$.*

Corollary 2.4 *If F has a cone decomposition with the compatibility Assumption 1 for $s = d-1$ and also $m = \text{Cat}(F)$, then $\text{Cat}(X) \leq \text{Cat}(F) + \lceil \frac{\dim B}{d} \rceil$.*

Remark 2.5 *Without Assumption 1, we only have*

$$\text{Cat}(X)+1 \leq (\text{Cat}(F)+1) \cdot (\text{Cat}(B)+1)$$

which is obtained immediately from the definition of Cat by Ganea [5] and the corresponding results of Varadarajan [31] and Hardie [8] for cat. For example, the principal bundle $\text{Sp}(1) \hookrightarrow \text{Sp}(2) \rightarrow S^7$ does satisfy Assumption 1 for $d \leq 3$, but not if $d \geq 4$, and we have $\text{Cat}(\text{Sp}(2)) \leq \text{Cat}(\text{Sp}(1)) + [\frac{7}{3}] = 3 > 2 = \text{Cat}(\text{Sp}(1)) + [\frac{7}{4}]$. In fact by Schweitzer [24], we know $\text{Cat}(\text{Sp}(2)) = 3$.

Remark 2.6 *By Remark 2.2 (2), Theorem 2.3 generalises Theorem 2.1.*

By applying this, we first obtain the following general result:

Theorem 2.7 *Let $C_m < \text{SU}(n)$ be a central (cyclic) subgroup of order m . Then we have $\text{Cat}(\text{SU}(n)/C_m) \leq 3(n-1)$ for all $n \geq 1$.*

This might be best possible, because we also obtain the following result.

Theorem 2.8 *We have*

$$\text{Cat}(\text{SU}(p^r)/C_{p^s}) = \text{cat}(\text{SU}(p^r)/C_{p^s}) = \text{rcat}(\text{SU}(p^r)/C_{p^s}) = 3(p^r - 1)$$

where p is a prime and $1 \leq s \leq r$.

Similarly we obtain the following result.

Theorem 2.9 *We have*

$$\begin{aligned} \text{Cat}(\text{SO}(6)) &= \text{cat}(\text{SO}(6)) = \text{cup}(\text{SO}(6)) = 9, \\ \text{Cat}(\text{SO}(7)) &= \text{cat}(\text{SO}(7)) = \text{cup}(\text{SO}(7)) = 11, \\ \text{Cat}(\text{SO}(8)) &= \text{cat}(\text{SO}(8)) = \text{cup}(\text{SO}(8)) = 12, \\ \text{Cat}(\text{SO}(9)) &= \text{cat}(\text{SO}(9)) = \text{cup}(\text{SO}(9)) = 20, \\ \text{Cat}(\text{PO}(8)) &= \text{cat}(\text{PO}(8)) = \text{cup}(\text{PO}(8)) = 18. \end{aligned}$$

Remark 2.10 *Theorem 2.3 also provides an alternative proof for a result of James-Singhof [16], that is, $\text{Cat}(\text{SO}(5)) = \text{cat}(\text{SO}(5)) = \text{cup}(\text{SO}(5)) = 8$ (see Section 4).*

We summarise all the known cases in the following table, where each number given in the right hand side of a connected, compact, simple Lie group indicates

its L-S category.

rank	1		2		3		4		$n (\geq 5)$	
A_n	SU(2)	1	SU(3)	2	SU(4)	3	SU(5)	4	SU($n+1$)	n
					SO(6)	9			$\vdots$	
	PU(2)	3	PU(3)	6	PU(4)	9	PU(5)	12	PU($n+1$)	—
B_n	Spin(3)	1	Spin(5)	3	Spin(7)	5	Spin(9)	—	Spin($2n+1$)	—
	SO(3)	3	SO(5)	8	SO(7)	11	SO(9)	20	SO($2n+1$)	—
C_n	Sp(1)	1	Sp(2)	3	Sp(3)	5	Sp(4)	—	Sp(n)	—
	PSp(1)	3	PSp(2)	8	PSp(3)	—	PSp(4)	—	PSp(n)	—
D_n					Spin(6)	3	Spin(8)	6	Spin($2n$)	—
					SO(6)	9	SO(8)	12	SO($2n$)	—
					PO(6)	9	PO(8)	18	PO($2n$)	—
									Ss($2n$)	—
Except. types			G_2	4			F_4	—	E_6, E_7, E_8	—

where "—" indicates the unknown case.

Remark 2.11 We recall that $A_1 = B_1 = C_1$, $B_2 = C_2$ and $A_3 = D_3$, and that the semi-spinor group $Ss(2n)$ is defined only for n even.

Taking into account the above table, we get the following by Theorem 1.6:

Corollary 2.12 The Ganea conjecture on L-S category holds for every connected, compact, simple Lie group G when L-S category is known as above.

The paper is organised as follows; In Section 3 we prove Theorem 2.3. In Section 4 we determine $\text{cat}(\text{SO}(n))$ for $n = 5, 6, 7, 8, 9$ and $\text{cat}(\text{PO}(8))$. In Section 5 we prove Theorem 2.7 and determine $\text{cat}(\text{SU}(p^r)/C_{p^s})$.

3 Proof of Theorem 2.3

Let B_i be the $(d \cdot i + s)$ -skeleton of B and $n = [\frac{\dim B}{d}]$ the biggest integer not exceeding $\frac{\dim B}{d}$. Then by Ganea [5], Theorem 2.1 implies that there are n cofibre sequences $A_i \xrightarrow{\lambda_i} B_{i-1} \hookrightarrow B_i$, $1 \leq i \leq n$ with $B_0 = \{*\}$, $B_n = B$. Note

that A_i is $(d \cdot i - 2)$ -connected and of dimension $(d \cdot i + s - 1)$. Hence we obtain

$$\begin{aligned} B_i &= B_{i-1} \cup_{\lambda_i} C(A_i), \quad \lambda_i : A_i \rightarrow B_{i-1} \\ A_i &= A_i^{(d \cdot i + s - 1)} = \bigcup_{a=0}^s A_i^{(d \cdot i + a - 1)}, \quad 1 \leq i \leq n, \\ B_0 &= \{*\}, \quad B_n \simeq B. \end{aligned}$$

Then there is a filtration of E by $E|_{B_i}$, $0 \leq i \leq n$, as follows (see Whitehead [32], for example):

$$\begin{aligned} E|_{B_i} &= E|_{B_{i-1}} \cup_{\Lambda_i} C(A_i) \times G, \quad \Lambda_i : A_i \times G \rightarrow E|_{B_{i-1}}, \quad 1 \leq i \leq n, \\ E|_{B_0} &= \{*\} \times G, \quad E|_{B_n} \simeq E, \end{aligned}$$

and $\tilde{\lambda}_i = \Lambda_i|_{A_i} : A_i \rightarrow E|_{B_{i-1}}$ gives a lift of $\lambda_i : A_i \rightarrow B_{i-1}$. Then by induction on i , we have

$$\begin{aligned} E|_{B_i} &= \{*\} \times G \cup_{\Lambda_1} C(A_1) \times G \cup_{\Lambda_2} \cdots \cup_{\Lambda_i} C(A_i) \times G, \\ \Lambda_i : A_i \times G &\xrightarrow{\tilde{\lambda}_i \times 1_G} E|_{B_{i-1}} \times G \\ &= \left(\{*\} \times G \cup_{\Lambda_1} C(A_1) \times G \cdots \cup_{\Lambda_{i-1}} C(A_{i-1}) \times G \right) \times G \\ &\xrightarrow{1 \times \mu} \{*\} \times G \cup_{\Lambda_1} C(A_1) \times G \cdots \cup_{\Lambda_{i-1}} C(A_{i-1}) \times G = E|_{B_{i-1}}, \end{aligned}$$

where μ is the multiplication of G . For dimensional reasons, we may regard

$$\tilde{\lambda}_i : (A_i, A_i^{(d \cdot i + a - 1)}) \rightarrow (E^{(d \cdot i + s - 1)}|_{B_{i-1}}, E^{(d \cdot i + a - 1)}|_{B_{i-1}}), \quad 0 \leq a \leq s,$$

and $\mu(G^{(i)} \times G^{(j)}) \subset G^{(i+j)}$ up to homotopy. Then we have the following descriptions for all $k \geq d \cdot i - 1$ and $j \geq d - 1$:

$$\begin{aligned} E^{(k)}|_{B_i} &= (\{*\} \times G \cup_{\Lambda_1} C(A_1) \times G \cup_{\Lambda_2} \cdots \cup_{\Lambda_i} C(A_i) \times G)^{(k)}, \\ &= \left(\begin{aligned} &\{*\} \times G^{(k)} \cup_{\Lambda_1} \bigcup_{\ell=0}^s (C(A_1^{(d \cdot 1 + \ell - 1)}) \times G^{(k - d - \ell)}) \\ &\cdots \cup_{\Lambda_i} \bigcup_{\ell=0}^s (C(A_i^{(d \cdot i + \ell - 1)}) \times G^{(k - d \cdot i - \ell)}) \end{aligned} \right), \\ \Lambda_i : A_i^{(d \cdot i + \ell - 1)} \times G^{(j - \ell)} &\xrightarrow{\tilde{\lambda}_i \times 1_{G^{(j)}}} E^{(d \cdot i + \ell - 1)}|_{B_{i-1}} \times G^{(j - \ell)} \\ &= \left(\begin{aligned} &\{*\} \times G^{(d \cdot i + \ell - 1)} \\ &\cup_{\Lambda_1} \bigcup_{a=0}^s (C(A_1^{(d \cdot 1 + a - 1)}) \times G^{(d \cdot (i-1) + \ell - a - 1)}) \\ &\cdots \cup_{\Lambda_{i-1}} \bigcup_{a=0}^s (C(A_{i-1}^{(d \cdot (i-1) + a - 1)}) \times G^{(d \cdot (i-1) + \ell - a - 1)}) \end{aligned} \right) \times G^{(j - \ell)} \\ &\xrightarrow{1 \times \mu} \left(\begin{aligned} &\{*\} \times G^{(d \cdot i + j - 1)} \cup_{\Lambda_1} \bigcup_{a=0}^s (C(A_1^{(d \cdot 1 + a - 1)}) \times G^{(d \cdot (i-1) + j - a - 1)}) \\ &\cdots \cup_{\Lambda_{i-1}} \bigcup_{a=0}^s (C(A_{i-1}^{(d \cdot (i-1) + a - 1)}) \times G^{(d \cdot (i-1) + j - a - 1)}) \end{aligned} \right) \end{aligned}$$

$$\begin{aligned}
&= \left(\{*\} \times G \cup_{\Lambda_1} C(A_1) \times G \cup_{\Lambda_2} \cdots \cup_{\Lambda_{i-1}} C(A_{i-1}) \times G \right)^{(d \cdot i + j - 1)} \\
&= E^{(d \cdot i + j - 1)}|_{B_{i-1}}.
\end{aligned}$$

Similarly, we obtain the following filtration $\{E'_k\}_{0 \leq k \leq n+m}$ of $E \times_G F$.

$$\begin{aligned}
E'_k &= \begin{cases} F_k \cup_{\Lambda'_1} C(A_1) \times F_{k-1} \cup_{\Lambda'_2} \cdots \cup_{\Lambda'_k} C(A_k) \times F_0, & k \leq n, \\ F_k \cup_{\Lambda'_1} C(A_1) \times F_{k-1} \cup_{\Lambda'_2} \cdots \cup_{\Lambda'_n} C(A_n) \times F_{k-n}, & n \leq k, \end{cases} \\
\Lambda'_i : A_i \times F_j &\xrightarrow{\tilde{\lambda}_i \times 1_{F_j}} E^{(d \cdot i + s - 1)}|_{B_{i-1}} \times F_j \\
&= \left(G^{(d \cdot i + s - 1)} \cup_{\Lambda_1} \bigcup_{a=0}^s (C(A_1^{(d+a-1)}) \times G^{(d \cdot (i-1) + s - a - 1)}) \right. \\
&\quad \left. \cdots \cup_{\Lambda_{i-1}} \bigcup_{a=0}^s (C(A_{i-1}^{(d \cdot (i-1) + a - 1)}) \times G^{(d \cdot s - a - 1)}) \right) \times F_j \\
&\xrightarrow{1 \times \psi} \left(F_{i+j-1} \cup_{\Lambda'_1} \bigcup_{a=0}^s (C(A_1^{(d+a-1)}) \times F_{i+j-2}) \right. \\
&\quad \left. \cdots \cup_{\Lambda'_{i-1}} \bigcup_{a=0}^s (C(A_{i-1}^{(d \cdot (i-1) + a - 1)}) \times F_j) \right) \\
&= F_{i+j-1} \cup_{\Lambda'_1} C(A_1) \times F_{i+j-2} \cdots \cup_{\Lambda'_{i-1}} C(A_{i-1}) \times F_j \\
&= E'_{i+j-1}|_{B_{i-1}},
\end{aligned}$$

since $\psi(G^{(d \cdot (\ell+1) + s - a - 1)} \times F_j) \subseteq \psi(G^{(d \cdot (\ell+1) + s - 1)} \times F_j) \subset F_{\ell+j}$ by Assumption 1. The above definition of Λ'_i also determines a map

$$\psi_{i,j} : E^{(d \cdot (i+1) + s - 1)}|_{B_i} \times F_j \longrightarrow E'_{i+j}|_{B_i}$$

so that $\Lambda'_i = \psi_{i-1,j} \circ (\tilde{\lambda}_i \times 1)$. Let us recall that $F_j = F_{j-1} \cup_{\rho_j} C(K_j)$ for $1 \leq j \leq m$. Then the definition of E'_k implies

$$E'_k = \begin{cases} E'_{k-1} \cup C(K_k) \cup C(A_1) \times C(K_{k-1}) \cup \cdots \\ \quad \cdots \cup C(A_{k-1}) \times C(K_1) \cup C(A_k) \times \{*\} & \text{for } k \leq n, \\ E'_{k-1} \cup C(K_k) \cup C(A_1) \times C(K_{k-1}) \cup \cdots \\ \quad \cdots \cup C(A_{n-1}) \times C(K_{k-n+1}) \cup C(A_n) \times C(K_{k-n}) & \text{for } k > n. \end{cases}$$

To observe the relation between $\text{Cat}(E'_{k-1})$ and $\text{Cat}(E'_k)$, we introduce the following two relative homeomorphisms:

$$\chi(\rho_j) : (C(K_j), K_j) \rightarrow (F_{j-1} \cup C(K_j), F_{j-1}) (= (F_j, F_{j-1}))$$

$$\begin{aligned}\chi(\tilde{\lambda}_i) : (C(A_i), A_i) &\rightarrow (E^{(d \cdot i + s - 1)}|_{B_i} \cup C(A_i), E^{(d \cdot i + s - 1)}|_{B_{i-1}}) \\ &(\subset (E^{(d \cdot i + s)}|_{B_i}, E^{(d \cdot i + s - 1)}|_{B_{i-1}})).\end{aligned}$$

Then the attaching map of $C(A_i) \times C(K_j)$ is given by the Whitehead product $[\chi(\tilde{\lambda}_i), \chi(\rho_j)] : A_i * K_j = (C(A_i) \times K_j) \cup (A_i \times C(K_j)) \rightarrow E'_{i+j-1}$ defined as follows:

$$\begin{aligned}[\chi(\tilde{\lambda}_i), \chi(\rho_j)]|_{C(A_i) \times K_j} : C(A_i) \times K_j &\xrightarrow{\chi(\tilde{\lambda}_i) \times 1} E^{(d \cdot i + s)}|_{B_i} \times F_{j-1} \\ &\subseteq E^{(d \cdot (i+1) + s - 1)}|_{B_i} \times F_{j-1} \xrightarrow{\psi_{i,j-1}} E'_{i+j-1}|_{B_i} \subseteq E'_{i+j-1}, \\ [\chi(\tilde{\lambda}_i), \chi(\rho_j)]|_{A_i \times C(K_j)} : A_i \times C(K_j) &\xrightarrow{\tilde{\lambda}_i \times \chi(\rho_j)} E^{(d \cdot i + s - 1)}|_{B_{i-1}} \times F_j \\ &\xrightarrow{\psi_{i-1,j}} E'_{i+j-1}|_{B_{i-1}} \subseteq E'_{i+j-1}.\end{aligned}$$

This implies immediately that $\text{Cat}(E'_k) \leq \text{Cat}(E'_{k-1}) + 1$. Then by induction on k , we obtain that $\text{Cat}(E'_k) \leq k$. Thus we have $\text{Cat}(X) = \text{Cat}(E \times_G F) = \text{Cat}(E'_{m+n}) \leq m+n \leq m + \frac{\dim B}{d}$. This completes the proof of Theorem 2.3.

4 Proof of Theorem 2.9

As is well known, we have the following principal bundles (see for example [2], [34] and [9] in particular for the last fibration):

$$\begin{aligned}\text{Sp}(1) &\longrightarrow \text{Sp}(2) \longrightarrow S^7, \\ \text{SU}(3) &\longrightarrow \text{SU}(4) \longrightarrow S^7, \\ G_2 &\longrightarrow \text{Spin}(7) \longrightarrow S^7, \\ \text{Spin}(7) &\longrightarrow \text{Spin}(9) \longrightarrow S^{15}, \\ G_2 &\longrightarrow \text{Spin}(8) \longrightarrow S^7 \times S^7.\end{aligned}$$

Each scalar matrix $(-1) \in \text{Sp}(2)$ and $(-1) \in \text{SU}(4)$ acts on S^7 as the antipodal map, and so does the center of $\text{Spin}(7)$. Similarly the center of $\text{Spin}(9)$ acts on S^{15} as the antipodal map. Recall that the center of $\text{Spin}(8)$ is isomorphic to $\mathbb{Z}/2 \times \mathbb{Z}/2$, each generator of which acts on S^7 as the antipodal map respectively. Since there are isomorphisms $\text{Sp}(2) \cong \text{Spin}(5)$ and $\text{SU}(4) \cong \text{Spin}(6)$, we obtain principal bundles:

$$\begin{aligned}\text{Sp}(1) &\longrightarrow \text{SO}(5) \longrightarrow \mathbb{R}P^7, \\ \text{SU}(3) &\longrightarrow \text{SO}(6) \longrightarrow \mathbb{R}P^7, \\ G_2 &\longrightarrow \text{SO}(7) \longrightarrow \mathbb{R}P^7, \\ \text{Spin}(7) &\longrightarrow \text{SO}(9) \longrightarrow \mathbb{R}P^{15}, \\ G_2 &\longrightarrow \text{PO}(8) \longrightarrow \mathbb{R}P^7 \times \mathbb{R}P^7.\end{aligned}$$

Cone decompositions of the fibres except $\text{Spin}(7)$ are given as follows (see Theorem 2.1 of [13] for G_2):

$$\begin{aligned} * &\subset \text{Sp}(1) = S^3, \\ * &\subset \text{SU}(3)^{(5)} \subset \text{SU}(3), \\ * &\subset G_2^{(5)} \subset G_2^{(8)} \subset G_2^{(11)} \subset G_2, \end{aligned}$$

where $\text{SU}(3)^{(5)} = G_2^{(5)} = \Sigma\mathbb{C}P^2$, $\text{SU}(3) = \text{SU}(3)^{(5)} \cup CS^7$, $G_2^{(8)} \simeq G_2^{(5)} \cup C(S^5 \cup e^7)$, $G_2^{(11)} \simeq G_2^{(8)} \cup C(S^8 \cup e^{10})$ and $G_2 = G_2^{(11)} \cup CS^{13}$. Since these fibres satisfy the conditions in Remark 2.2 (1), we obtain $\text{Cat}(\text{SO}(5)) \leq 8$, $\text{Cat}(\text{SO}(6)) \leq 9$, $\text{Cat}(\text{SO}(7)) \leq 11$ and $\text{Cat}(\text{PO}(8)) \leq 18$ using Theorem 2.3. By virtue of the mod 2 cup-lengths we have that $\text{cup}(\text{SO}(5)) \geq 8$, $\text{cup}(\text{SO}(6)) \geq 9$, $\text{cup}(\text{SO}(7)) \geq 11$ and $\text{cup}(\text{PO}(8)) \geq 18$ respectively. Thus we obtain the results for $\text{SO}(5)$, $\text{SO}(6)$, $\text{SO}(7)$ and $\text{PO}(8)$.

A cone decomposition of $\text{Spin}(7)$ is given as follows in [14]:

$$* = F_0 \subset F_1 \subset F_2 \subset F_3 \subset F_4 \subset F_5 = \text{Spin}(7),$$

where $F_1 = \text{SU}(4)^{(7)}$, $F_2 = \text{SU}(4)^{(12)} \cup e^6$, $F_3 = \text{SU}(4) \cup e^6 \cup e^9 \cup e^{11} \cup e^{13}$ and $F_4 = \text{Spin}(7)^{(18)}$. We need here to check if the filtration satisfies Assumption 1; the only problem is to determine whether $\psi|_{\text{Spin}(7)^{(3)} \times F_1} : \text{Spin}(7)^{(3)} \times F_1 \rightarrow F$ is compressible into F_4 or not. Since $\text{Spin}(7)^{(3)}$ and F_1 are included in $\text{SU}(4) \subset F_4$, we have $\text{Im}(\psi|_{\text{Spin}(7)^{(3)} \times F_1}) \subset F_4$. Then we obtain $\text{Cat}(\text{SO}(9)) \leq 20$ using Theorem 2.3. The mod 2 cup-length implies that $\text{cup}(\text{SO}(9)) \geq 20$. Thus we obtain the result for $\text{SO}(9)$.

Since $\text{SO}(8)$ is homeomorphic to $\text{SO}(7) \times S^7$, we easily see that

$$\text{Cat}(\text{SO}(8)) \leq \text{Cat}(\text{SO}(7)) + \text{Cat}(S^7) = 12$$

by Takens [?]. The mod 2 cup-length implies that $\text{cup}(\text{SO}(8)) \geq 12$. Thus we obtain the result for $\text{SO}(8)$. This completes the proof of Theorem 2.9.

5 Proof of Theorems 2.7 and 2.8

Firstly, we show Theorem 2.7. The following principal bundle is well-known:

$$\text{SU}(n-1) \longrightarrow \text{SU}(n) \longrightarrow S^{2n-1}.$$

The central (cyclic) subgroup C_m of $\text{SU}(n)$ acts on S^{2n-1} freely and hence we obtain a principal bundle:

$$\text{SU}(n-1) \longrightarrow \text{SU}(n)/C_m \longrightarrow L^{2n-1}(m),$$

where $L^{2n-1}(m)$ is a lens space of dimension $2n-1$.

A cone decomposition of $SU(n-1)$ is constructed by Kadzisa [17]:

$$* \subset V \subset V^2 \subset \cdots \subset V^{n-2} = SU(n-1),$$

where $V^k \subseteq SU(n-1)$ is a representing subspace of the quotient module $H^*(SU(n-1))/D^{k+1}$ and D^{k+1} is the submodule generated by products of $k+1$ elements in positive degrees, which satisfies $V^i \cdot V^j \subseteq V^{i+j}$ for any i and j . Thus V is the subcomplex $S^3 \cup e^5 \cup e^7 \cup \cdots \cup e^{2n-3}$ of $SU(n-1)$ which is homeomorphic to ΣCP^{n-2} (see [33], for example). Then Assumption 1 is automatically satisfied, and hence using $SU(n-1)^{(k)} \subset V^k$, we obtain

$$\text{Cat}(SU(n)/C_m) \leq 3(n-1)$$

by Theorem 2.3. This completes the proof of Theorem 2.7.

Secondly, we show Theorem 2.8. By Rudyak [20] [21] and Strom [30], we know the following proposition.

Proposition 5.1 (Rudyak [20] [21], Strom [30]) *Let h be a cohomology theory. For an element $u \in \tilde{h}^*(X)$, let $\text{wgt}(u; h)$ be the minimal number k such that $(e_k^X)^*(u) \neq 0$ where $e_k^X : P^k \Omega X \rightarrow P^\infty \Omega X \simeq X$, which satisfies*

- (1) *We have $\text{wgt}(0; h) = \infty$ and $\infty > \text{wgt}(u; h) \geq 1$ for any $u \neq 0$ in $\tilde{h}^*(X)$.*
- (2) *For any cohomology theory h , we have*

$$\min \{ \text{wgt}(u; h), \text{wgt}(v; h) \} \leq \text{wgt}(u + v; h).$$

- (3) *For any multiplicative cohomology theory h , we have*

$$\text{wgt}(u; h) + \text{wgt}(v; h) \leq \text{wgt}(u \cdot v; h).$$

- (4) $\text{wgt}(X; h) = \max \{ \text{wgt}(u; h) \mid u \in \tilde{h}^*(X), u \neq 0 \}.$

Let us recall that, for any compact Lie group G , the ordinary cohomology of ΩG is concentrated in even degrees. Then, for any element u of even degree in $\tilde{H}^*(G; \mathbb{Z}/p)$, we have $\text{wgt}(u; H\mathbb{Z}/p) \geq 2$, since $P^1(\Omega G) = \Sigma \Omega(G)$.

The cohomology rings of $SU(p^r)/C_{p^s}$ for a prime p and $1 \leq s \leq r$ are given as follows (see [3]):

$$H^*(SU(p^r)/C_{p^s}; \mathbb{Z}/p) = \mathbb{Z}/p[x_2]/(x_2^{p^r}) \otimes \wedge(x_1, x_3, \dots, x_{2p^r-3}).$$

Note that $x_1^2 = x_2$ if $p = 2$ and $s = 1$. Then, using Proposition 5.1, we obtain

$$\text{wgt}(SU(p^r)/C_{p^s}; H\mathbb{Z}/p) \geq \text{wgt}(x_1 \cdot x_2^{p^r-1} \cdot x_3 \cdots x_{2p^r-3}; H\mathbb{Z}/p) \geq 3(p^r - 1),$$

since $\text{wgt}(x_2; H\mathbb{Z}/p) \geq 2$. Thus we have the following lemma.

Lemma 5.2 $\text{rcat}(\text{SU}(p^r)/C_{p^s}) \geq 3(p^r - 1)$ for any prime p and $1 \leq s \leq r$.

By using Theorem 2.7, we obtain Theorem 2.8.

References

- [1] J. F. Adams, “Stable homotopy and generalized homology”, Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1974.
- [2] J. F. Adams, “Lectures on exceptional Lie groups” (with a foreword by J. Peter May, edited by Zafer Mahmud and Mamoru Mimura) Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1996.
- [3] P. F. Baum and W. Browder, *The cohomology of quotients of classical groups*, *Topology* **3** (1965), 305–336.
- [4] L. Fernández-Suárez, A. Gómez-Tato, J. Strom, D. Tanré, *The Lusternik-Schnirelmann category of $\text{Sp}(3)$* , *Trans. Amer. Math. Soc.* **132** (2004), 587–595 (electronic).
- [5] T. Ganea, *Lusternik-Schnirelmann category and strong category*, *Illinois J. Math.* **11** (1967), 417–427.
- [6] T. Ganea, *Some problems on numerical homotopy invariants*, In: “Symposium on Algebraic Topology”, 13–22, *Lect. Notes in Math.* **249**, Springer Verlag, Berlin 1971.
- [7] J. C. Gómez-Larrañaga and F. González-Acuña, *Lusternik-Schnirelmann category of 3-manifolds*, *Topology* **31** (1992), 791–800.
- [8] K. A. Hardie, *A note on fibrations and category*, *Michigan Math. J.* **17**, (1970), 351–352.
- [9] F. R. Harvey, “Spinors and calibrations”, *Perspectives in Mathematics* **9**, Academic Press, Boston, 1990.
- [10] N. Iwase, *Ganea’s conjecture on Lusternik-Schnirelmann category*, *Bull. Lon. Math. Soc.* **30** (1998), 623–634.
- [11] N. Iwase, *A_∞ -method in Lusternik-Schnirelmann category*, *Topology* **41** (2002), 695–723.
- [12] N. Iwase, *Lusternik-Schnirelmann category of a sphere-bundle over a sphere*, *Topology* **42** (2003), 701–713.
- [13] N. Iwase, M. Mimura, *L-S categories of simply-connected compact simple Lie groups of low rank*, In: “Algebraic Topology: Categorical Decomposition Techniques”, (Isle of Skye, 2001), 199–212, *Progr. Math.*, 215, Birkhäuser Verlag, Basel, 2004.

- [14] N. Iwase, M. Mimura, T. Nishimoto, *On the cellular decomposition and the Lusternik-Schnirelmann category of $\text{Spin}(7)$* , Topology Appl. **133** (2003), 1–14.
- [15] I. M. James, *On category, in the sense of Lusternik-Schnirelmann*, Topology **17** (1978), 331–348.
- [16] I. M. James, W. Singhof, *On the category of fibre bundles, Lie groups, and Frobenius maps*, Higher Homotopy Structures in Topology and Mathematical Physics (Poughkeepsie, NY, 1996), 177–189, Contemp. Math. **227**, Amer. Math. Soc., Providence, 1999.
- [17] H. Kadzisa, *Cone-Decompositions of the Special Unitary Groups*, Topology Appl., to appear.
- [18] L. Montejano, *A quick proof of Singhof's $\text{cat}(M \times S^1) = \text{cat}(M) + 1$ theorem*, Manuscripta Math. **42** (1983), 49–52.
- [19] J. Oprea, Y. Rudyak, *Detecting elements and Lusternik-Schnirelmann category of 3-manifolds*, In: “Lusternik-Schnirelmann category and related topics” (South Hadley, MA, 2001), 181–191, Contemp. Math. **316**, Amer. Math. Soc., Providence, 2002.
- [20] Y. B. Rudyak, *On the Ganea conjecture for manifolds*, Proc. Amer. Math. Soc. **125** (1997), 2511–2512.
- [21] Y. B. Rudyak, *On category weight and its applications*, Topology **38** (1999), 37–55.
- [22] Y. B. Rudyak, *On analytical applications of stable homotopy (the Arnold conjecture, critical points)*, Math. Z. **230** (1999), 659–672.
- [23] M. Sakai, *A proof of the homotopy push-out and pull-back lemma*, Proc. Amer. Math. Soc. **129** (2001), 2461–2466.
- [24] P. A. Schweitzer, *Secondary cohomology operations induced by the diagonal mapping*, Topology **3** (1965), 337–355.
- [25] W. Singhof, *On the Lusternik-Schnirelmann category of Lie groups*, Math. Z. **145** (1975), 111–116.
- [26] W. Singhof, *On the Lusternik-Schnirelmann category of Lie groups, II*, Math. Z. **151** (1976), 143–148.
- [27] W. Singhof, *Generalized higher order cohomology operations induced by the diagonal mapping*, Math. Z. **162** (1978), 7–26.
- [28] W. Singhof, *Minimal coverings of manifolds with balls*, Manuscripta Math. **29** (1979), 385–415.
- [29] J. D. Stasheff, *Homotopy associativity of H-spaces, I, II*, Trans. Amer. Math. Soc. **108** (1963), 275–292, 293–312.
- [30] J. Strom, *Essential category weight and phantom maps*, in: Cohomological methods in homotopy theory (Bellaterra, 1998), 409–415, Progr. Math. **196**, Birkhaeuser, Basel, 2001.

- [31] K. Varadarajan, *On fibrations and category*, Math. Z. **88** (1965), 267–273.
- [32] G. W. Whitehead, “Elements of Homotopy Theory”, Graduate Texts in Mathematics **61**, Springer Verlag, Berlin, 1978.
- [33] I. Yokota, *On the cell structures of $SU(n)$ and $Sp(n)$* , Proc. Japan Acad. **31** (1955), 673–677.
- [34] I. Yokota, “Groups and Representations” (Japanese), Shōkabō, Tokyo, 1973.